

Appendix 3

Indicators where Nottinghamshire is either worse than England or the trend is getting worse.

Overarching indicators

- Healthy life expectancy at birth | Female | All ages
- Life expectancy at birth | Female | All ages
- Disability free life expectancy at birth | Female | All ages
- Healthy life expectancy at 65 | Male | 65
- Disability-free life expectancy at 65 | Female | 65

Wider determinants of health

- Children in absolute low income families (under 16s) | Persons | <16 yrs
- Children in relative low income families (under 16s) | Persons | <16 yrs
- 16 to 17 year olds not in education, employment or training (NEET) | Persons | 16-17 yrs
- Adults with a learning disability who live in stable and appropriate accommodation | Persons | 18-64 yrs
- Adults in contact with secondary mental health services who live in stable and appropriate accommodation | Persons | 18-69 yrs
- The percentage of the population who are in receipt of long term support for a learning disability that are in paid employment | Persons | 18-64 yrs
- The percentage of the population who are in contact with secondary mental health services and on the Care Plan Approach, that are in paid employment | Persons | 18-69 yrs
- Sickness absence: percentage of working days lost due to sickness absence | Persons | 16+ yrs
- Social Isolation: percentage of adult carers who have as much social contact as they would like | Persons | All ages

Health improvement

- Obesity in early pregnancy | Female | Not applicable
- Smoking in early pregnancy | Female | Not applicable
- Baby's first feed breastmilk | Persons | Newborn
- Breastfeeding prevalence at 6-8 weeks after birth - current method | Persons | 6-8 weeks
- Smoking status at time of delivery | Female | All ages
- Year 6: Prevalence of overweight (including obesity) | Persons | 10-11 yrs
- Hospital admissions caused by unintentional and deliberate injuries in young people | Persons | 15-24 yrs
- Percentage of adults classified as overweight or obese | Persons | 18+ yrs
- Successful completion of drug treatment: opiate users | Persons | 18+ yrs
- Successful completion of drug treatment: non opiate users | Persons | 18+ yrs
- Successful completion of alcohol treatment | Persons | 18+ yrs
- Adults with substance misuse treatment need who successfully engage in community based structured treatment following release from prison | Persons | 18+ yrs
- Admission episodes for alcohol-related conditions (Narrow) | Persons | All ages
- Percentage of cancers diagnosed at stages 1 and 2 | Persons | All ages
- Cancer screening coverage: breast cancer | Female | 53-70 yrs
- Cancer screening coverage: cervical cancer | Female | 50-64 yrs
- Newborn Hearing Screening: Coverage | Persons | <1 yr
- Newborn and Infant Physical Examination Screening Coverage | Persons | <1 yr
- Cumulative percentage of the eligible population offered an NHS Health Check | Persons | 40-74 yrs
- Percentage reporting a long-term Musculoskeletal (MSK) problem | Persons | 16+ yrs

Health protection

- Population vaccination coverage: Dtap IPV Hib (1 year old) | Persons | 1 yr
- Population vaccination coverage: Dtap IPV Hib | Persons | 2 yrs
- Population vaccination coverage: PCV booster | Persons | 2 yrs
- Population vaccination coverage: Hib and MenC booster | Persons | 2 yrs
- Population vaccination coverage: MMR for one dose | Persons | 5 yrs
- Population vaccination coverage: HPV vaccination coverage for one dose | Male | 12-13 yrs
- Population vaccination coverage: HPV vaccination coverage for one dose | Female | 12-13 yrs
- Population vaccination coverage: HPV vaccination coverage for two doses | Female | 13-14 yrs
- Population vaccination coverage: Meningococcal ACWY conjugate vaccine (MenACWY) | Persons | 14-15 yrs
- Adjusted antibiotic prescribing in primary care by the NHS | Persons | All ages

Healthcare and early deaths

- Under 75 mortality rate from cancer | Persons | <75 yrs
- Excess under 75 mortality rate in adults with severe mental illness (SMI) | Persons | 18-74 yrs
- Suicide rate | Persons | 10+ yrs
- Preventable sight loss: age related macular degeneration (AMD) | Persons | 65+ yrs
- Preventable sight loss: sight loss certifications | Persons | All ages
- Hip fractures in people aged 80 and over | Persons | 80+ yrs